

VOLINACTM

Diclofenac Sodium

PAIN
RELIEF
GEL



COMPOSITION

VOLINACTM Gel: Each gram gel contains Diclofenac Diethylamine BP equivalent to Diclofenac Sodium BP 10 mg.

PHARMACOLOGY

Diclofenac Diethylamine is systemically absorbed through the skin; it inhibits the enzyme cyclooxygenase, thus reducing the formation of PGE₂. Moreover, it also increases the uptake of Arachidonic acid into the cellular pool. Menthol is a vasodilator. It dilates the blood vessels, produces a feeling of coolness and produces analgesia. Methyl salicylate is a known anti-inflammatory agent.

INDICATION

Indicated for the quick relief from pain, swelling and inflammation due to musculo-skeletal disorders such as sprains, strains, tendinitis, bursitis, hands, neck & shoulder pain, sciatica, muscle stiffness, joint pain, back ache and lumbago.

DOSAGE AND ADMINISTRATION

Approximately one-inch band of gel should be applied to the affected site three to four times daily with rubbing till the film disappears.

CONTRAINDICATIONS

Known hypersensitivity to any part of the preparation.

PRECAUTIONS

For external use only. Avoid contact with the eyes. Stop use and ask a doctor if condition worsens or does not improve within 7 days. Keep out of the reach of children. If swallowed, get medical help or contact a poison control center right away.

SIDE-EFFECTS

Usually well tolerated. Extremely low frequency of hypersensitivity reactions.

USE IN PREGNANCY AND LACTATION

The safety of **VOLINACTM Gel** has not been established during pregnancy. There are no well-controlled studies of diclofenac in pregnant women.

USE IN PEDIATRIC PATIENTS

Safety and effectiveness in pediatric patients have not been established.

STORAGE

Store in a cool and dry place, protected from light.

HOW SUPPLIED

VOLINACTM Gel: Each box contains 50 gm tube.

Manufactured by



**SQUARE
PHARMACEUTICALS LTD.**

Pabna, Bangladesh

TM- Trade Mark

PAIN
RELIEF
GEL

