

LNCTM

Cilnidipine INN



Composition

LNCTM 5: Each film coated tablet contains Cilnidipine INN 5 mg.

LNCTM 10: Each film coated tablet contains Cilnidipine INN 10 mg.

Pharmacology

Cilnidipine binds to the dihydropyridine binding sites of the L-type voltage dependent calcium channel and inhibits Ca²⁺ influx across the cell membranes of vascular smooth muscle cells via this channel, consequently vascular smooth muscle is relaxed, causing vasodilation.

Cilnidipine inhibits Ca²⁺ influx via N-type voltage dependent calcium channels in the sympathetic nerve cell membrane. Consequently, release of norepinephrine from sympathetic nerve terminals are inhibited. Cilnidipine is considered to suppress the reflex increase in heart rate after blood pressure reduction.

Indication

Hypertension.

Dosage and Administration

Adults: 5-10 mg once daily after breakfast. Maximum dose: 20 mg once daily.

The safety of Cilnidipine in pediatric patients has not been established.

Use in the elderly: Since the elderly may be more susceptible to hypotension, therapy should be initiated with the lowest possible dose (5 mg).

Contraindication

Cilnidipine is contraindicated in patients with known sensitivity to Cilnidipine or any of the excipients or patients having cardiogenic shock, recent MI or acute unstable angina and severe aortic stenosis.

Side Effects

The most common side effects of Cilnidipine are: Dizziness; flushing; headache; hypotension; peripheral edema; palpitations; GI disturbances; increased micturition frequency; lethargy; eye pain; depression.

Precaution

Cilnidipine should be administered with care in the following patients: patients with serious hepatic dysfunction, patients with a history of serious adverse reactions to calcium antagonists. During the discontinuation, the dosage should be gradually decreased under close observation.

Pregnancy & Lactation

Cilnidipine should not be administered in pregnant woman or woman having possibilities of being pregnant. It is also advisable to avoid the administration of Cilnidipine to nursing mothers. However, if the administration is indispensable, the patient should be instructed to discontinue lactation.

Drug Interaction

Other anti-hypertensives & antipsychotics that cause hypotension, quinidine, carbamazepine, rifampicin, cimetidine, erythromycin.

Storage

Store below 30^o C, protect from light and moisture. Keep out of reach of children.

How Supplied

LNCTM 5: Each box contains 30 tablets in blister pack.

LNCTM 10: Each box contains 30 tablets in blister pack.

Manufactured by



SQUARE

PHARMACEUTICALS LTD.
BANGLADESH