

Bilberry 160 mg capsule

Ocubil™ 160



Natural aid for better eye health

COMPOSITION

Ocubil™ 160 capsule : Each capsule contains 160 mg standardized extract of Bilberry fruit (*Vaccinium myrtillus L.*).

DESCRIPTION

Bilberry is the name of a small european blueberry. The standardized extract of bilberry has become popular in the United States as dietary supplements over the past decade. Bilberry preparations consist of the whole, dried, ripe, black or bluish – black fruit of *Vaccinium myrtillus L.* Some concentrated extracts are standardized to anthocyanosides, calculated as 25% anthocyanidins, but may actually contain about 37% by weight. The medicinal parts are the dried leaves, the ripe, dried fruit and the ripe fresh fruit. The plant is common to Central and Northern Europe, Asia and North America. The bilberry contains tannins, anthocyanins and flavonoid glycosides.

PHARMACOLOGY

Bilberry extract is used to treat Retinopathy, Cataracts, Macular degeneration, Retinitis pigmentosa, Hemorrhagic retinopathy. Bilberry anthocyanins regenerate rhodopsin are indicated in the treatment of poor night vision and cataracts. Bilberry is also used to treat microcirculatory disorders which include varicose veins, atherosclerosis, venous insufficiency.

MECHANISM OF ACTION

- Inhibits enzymatic cleavage of collagen by the enzymes secreted by leucocytes during Inflammation.
- By inhibiting retinal phosphoglucomutase and glucose-6-phosphatase it confers retinal protection.
- Reduces microvascular impairments due to ischemic reperfusion injury, with preservation of endothelium, attenuation of leukocyte adhesion and improvement of capillary perfusion.
- Produces dose dependent inhibition of platelet aggregation and clot retraction.
- Decreases basement membrane collagen hydrolysis by significantly reducing
- permeability of the blood-brain barrier and increases recovery rate of the BBB caused by permeability-increasing agents.

INDICATION & USES

Retinopathy (hypertensive and diabetic), Night blindness, Cataracts, Macular degeneration, Retinitis pigmentosa and Hemorrhagic retinopathy.

DOSAGE & ADMINISTRATION

One **Ocubil™** 160 mg capsule should be taken 2 - 3 times daily or as per the instruction of a physician.

SIDE EFFECT

None known.

CONTRAINDICATION

None known.

PREGNANCY & LACTATION

No known restriction.

DRUG INTERACTION

None known. Very high doses (more than 170 mg anthocyanins daily for 30 - 60 days) may interact with warfarin or other antiplatelet drugs.

STORAGE

Store at cool and dry place below 30° c temperature away from direct sunlight & moisture. Keep the medicine out of the reach of children.

HOW SUPPLIED

Ocubil™ 160 : Each commercial box contains 30 capsules in Alu-PVDC blister pack.

Manufactured by

SQUARE PHARMACEUTICALS LTD.

HERBAL DIVISION

BSCIC, Pabna, Bangladesh

www.squarepharma.com.bd

