

Iventi-DTM Eye Drops

Moxifloxacin 0.5% &
Dexamethasone Phosphate 0.1%
Sterile Solution

Iventi-DTM Eye Drops is a combination of a fourth generation fluoroquinolone Moxifloxacin and a potent corticosteroid Dexamethasone. Moxifloxacin controls infection by inhibiting the DNA gyrase & Topoisomerase IV. Dexamethasone effectively controls the inflammation by inhibiting the release inflammatory mediators.

COMPOSITION

Iventi-DTM Eye Drops: Each ml contains Moxifloxacin 5 mg as Moxifloxacin Hydrochloride BP & Dexamethasone Phosphate 1 mg as Dexamethasone Sodium Phosphate BP.

INDICATIONS & USES

This combination is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infection exists. The combination can also be used for post-operative inflammation and any other ocular inflammation associated with infection.

DOSAGE & ADMINISTRATION

Prevention of post-operative ocular inflammation and infection: Instill 1 drop 4 times/day in the eye to be operated, starting 1 day before the surgery and during 15 days after the surgery.

Treatment of ocular infections caused by susceptible organisms: Instill 1 drop 4 times/day during 7 days, or as directed by the doctor.

CONTRAINDICATION

This ophthalmic solution is contraindicated in epithelial herpes simplex keratitis (Dendritic keratitis), vaccinia, varicella, and in many other viral diseases of the conjunctiva and cornea, Mycobacterial infection of the eye and fungal diseases of ocular structures and in individuals hypersensitive to any of the components of the medication.

PRECAUTION & WARNING

Prolonged use of steroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, and posterior subcapsular cataract formation.

USE IN PREGNANCY & LACTATION

There are no adequate and well-controlled studies in pregnant women. This drug should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Caution should be exercised when this drug is administered to a nursing woman.

USE IN CHILDREN

Safety and efficacy in pediatric patients have not been established.

SIDE EFFECTS

The most frequently reported drug-related undesirable effects seen with Moxifloxacin are conjunctival irritation, increased lacrimation, keratitis and papillary conjunctivitis.

STORAGE

Store in a cool & dry place, below 30°C, protected from light. Once the container is opened the contents must be used within 28 days. Discard after the 28 days. Keep out of reach of children.

HOW SUPPLIED

Iventi-DTM Eye Drops: Each LDPE container contains 5 ml of Moxifloxacin 0.5% and Dexamethasone Phosphate 0.1% of sterile solution.

Manufactured by



SQUARE

**PHARMACEUTICALS LTD.
BANGLADESH**