

Gynepro[®]

Vaginal Suppository

Metronidazole BP, Neomycin Sulphate BP, Polymyxin B Sulphate BP, Nystatin BP

COMPOSITION

Each **Gynepro[®]** vaginal suppository contains Metronidazole BP 200 mg, Neomycin Sulphate BP 35,000 IU, Polymyxin B Sulphate BP 35,000 IU, Nystatin BP 1,00,000 IU.

PHARMACOLOGY

Metronidazole, a nitroimidazole, is active against various anerobic bacteria and protozoa. Nystatin is a polyene antibiotic with antifungal activity. Neomycin is an aminoglycoside. It is bactericidal and active against some gram-positive and gram-negative organisms. Polymyxin is active against gram-negative organisms including *Pseudomonas aeruginosa*.

INDICATION

Vaginal trichomoniasis, vaginal leucorrhoeas, mixed vaginal infections (Fungal or Bacterial).

DOSAGE & ADMINISTRATION

One suppository in vagina at bedtime for 12 days, or as directed by the physician.

CONTRAINDICATION

Contraindicated to the patients who are hypersensitive to Metronidazole, Neomycin Sulphate, Polymyxin B Sulphate and Nystatin.

SIDE EFFECT

Skin rash, urticaria may occur rarely.

USE IN PREGNANCY AND LACTATION

Caution should be practiced in pregnancy and lactation.

PRECAUTION

Caution should be taken in case of renal impairment.

STORAGE

Store below 25° C temperature, protected from light & moisture.

HOW SUPPLIED

Gynepro[®]: Each box contains 12 suppositories in blister pack.

Manufactured by



SQUARE
PHARMACEUTICALS LTD.

Salgaria, Pabna, Bangladesh

® Registered Trade Mark