

Betameson-N[®]

Betamethasone and Neomycin Sulphate

Betameson-N[®] is very active fluorinated corticosteroid for topical use. At normal dosage, even often prolong use, **Betameson-N[®]** does not depress the suprarenal glands and one may consider its systemic activity as practically nil.

COMPOSITION

Each gram cream contains Betamethasone 1 mg as Betamethasone Dipropionate BP and Neomycin Sulphate BP 5 mg.

INDICATION

Severe inflammatory skin disorders in patients unresponsive to less potent corticosteroids.

These disorders include:

Atopic eczema, contact dermatitis, dyshidrosis, exfoliative dermatitis, infantile eczema, intertrigo, neurodermatitis, pruritus ani, pruritus vulve, psoriasis, seborrheic dermatitis, solar dermatitis.

DOSAGE

Apply sparingly 2-3 times daily, reducing frequency as condition

responds.

CONTRAINDICATION

Untreated bacterial, fungal or viral skin lesions, acne and perioral dermatitis.

PRECAUTION

Long-term continuous topical therapy should be avoided where possible, particularly in infants and children. If applied to the eyelids, care is needed to ensure that the preparation does not enter the eye, as glaucoma might occur.

SIDE EFFECT

Skin: burning, itching, irritation, dryness, folliculitis, hypopigmentation, perioral dermatitis, allergic contact dermatitis, hypertrichosis, acneiform eruptions.

With occlusive dressings: maceration of skin, secondary infection, atrophy, striae, miliaria.

HOW SUPPLIED

Each pack has a laminated tube containing 15 gm cream.

Manufactured by



SQUARE
PHARMACEUTICALS LTD.
BANGLADESH