# **Ceevit**® Forte

Vitamin C (Ascorbic acid)

Active Ingredient
Vitamin C (Ascorbic acid).

#### Indication

Treatment or prevention of Vitamin C Deficiency, Scurvy, Infection, Trauma, Burns, Cold exposure, Following Surgery, Fever, Stress, Cancer, Methaemoglobinaemia & Children receiving unfortified formulas. Also indicated in Hematuria, Dental Caries, Gum Diseases, Pyorrhea, Acne, Infertility, Atherosclerosis, Fractures, Leg ulcers, Hay fever, Vascular thrombosis prevention. Levodopa toxicity. Arsenic toxicity & etc.

### Dosage & Administration

One tablet daily with a meal or as directed by physician. Dissolve one tablet in half glass (100 ml) of water & drink instantly.

### Contraindication & Precaution

There is no serious contraindication to the administration of Vitamin C. Diabetes, patients prone to recurrent renal calculi & those on sodium restricted diets or anticoagulant therapy should not take excessive doses of ascorbic acid over an extended period of time.

#### Side Effect

Generally ascorbic acid is well tolerated. However, few side effects including stomach upset, diarrhea, mouth sores or frequent urination may be seen.

## Drug Interaction

Limited evidence suggests that ascorbic acid may influence the intensity & duration of action of by hydroxycoumarin.

Use in Pregnancy & Lactation

During pregnancy & lactation usual dose is safe.

Preparation

Vitamin C 1000 mg Effervescent Tablet.

Manufactured by

