

Urinary Alkalinising Agent

Uriset™

Potassium Citrate BP & Citric Acid Monohydrate BP

COMPOSITION

Uriset™ Oral Solution: Each 5 ml solution contains Potassium Citrate BP 1,500 mg and Citric Acid Monohydrate BP 250 mg.

PHARMACOLOGY

Uriset™ is a combined preparation of Potassium Citrate & Citric Acid Monohydrate. Potassium Citrate is absorbed and metabolized to Potassium Bicarbonate, thus acting as a systemic alkalinizer. The effects are essentially those of chlorides before absorption and those of bicarbonates subsequently. Oxidation is virtually complete so that less than 5% of the Potassium Citrate is excreted in the urine unchanged. It alkalinizes the urine without producing a systemic alkalosis in recommended dosage.

INDICATION

Uriset™ is indicated in following cases:

- To relieve discomfort in urinary tract infections due to decreased pH or acidic urine.
- In preventing kidney stone formation specially caused by uric acid with or without the presence of calcium stones.
- In gout therapy as a valuable adjuvant with uricosuric agents.
- Also effective in correcting acidosis caused by kidney disease.

DOSAGE AND ADMINISTRATION

Uriset™ Oral Solution should be taken diluted in water according to directions, followed by additional water, if desired. Palatability is enhanced if chilled before taking. Proper dilution may help prevent gastrointestinal injury associated with the oral ingestion of concentrated potassium salt preparations.

To relieve discomfort in UTI:

Adults and children over 6 years: 10 ml 3 times daily, diluted with 1 glass of water.

Children 1-6 years: 5 ml 3 times daily, diluted with ½ glass of water.

To prevent kidney stones & acidosis caused by kidney diseases:

Adults: 10-15 ml 4 times daily (or as directed by the physician) diluted with 1 glass of water.

Children: 5-10 ml 4 times daily (or as directed by the physician) diluted with ½ glass of water.

In gout therapy (with uricosuric agent):

Adults: 10-15 ml 4 times daily (or as directed by the physician), diluted with 1 glass of water.

Children: 5-10 ml 4 times daily (or as directed by the physician), diluted with ½ glass of water.

CONTRAINDICATION

Severe renal impairment with oliguria or azotemia, untreated Addison's disease, adynamia episodica hereditaria, acute dehydration, heat cramps, anuria, severe myocardial damage, and hyperkalemia from any cause. Known hypersensitivity to any ingredient.

SIDE EFFECT

This oral solution is generally well tolerated without any side effects when given in recommended doses to patients with normal renal function and urinary output.

WARNING & PRECAUTION

Large doses may cause hyperkalemia & alkalosis, especially in the presence of renal disease. Do not exceed recommended dosage. Discontinue use if adverse reaction occurs. Should be used with caution by patients with low urinary output unless under the supervision of a physician. As with all liquids containing a high concentration of potassium, patients should be directed to dilute adequately with water to minimize the possibility of gastrointestinal injury associated with the oral ingestion of concentrated potassium salt preparations; and preferable, to take each dose after meals to avoid saline laxative effect.

DRUG INTERACTION

Some products that may interact with this drug include: antacids that contain aluminum, aspirin and other salicylates (e.g., salsalate), certain blood pressure medications (e.g., ACE inhibitors such as lisinopril, angiotensin blockers such as losartan), drospirenone, eplerenone, certain heart medications (e.g., quinidine, digoxin), lithium, potassium supplements, certain "water pills" (potassium-sparing diuretics such as amiloride, spironolactone, triamterene).

PREGNANCY & LACTATION

Combination of Potassium Citrate & Citric Acid Monohydrate should not be used during pregnancy unless the potential benefit justifies the potential risk to the foetus.

It is not known if this drug is found in breast milk or not. Breast-feeding women should consult with their health care provider to confirm about taking this drug.

STORAGE CONDITION

Store below 30° C, protected from light. Keep out of the reach of children.

HOW SUPPLIED

Uriset™ Oral Solution: Each pack contains 200 ml of oral solution in PET bottle and a measuring cup.

Manufactured by



SQUARE
PHARMACEUTICALS LTD.
BANGLADESH