

Afun[®] VT

Clotrimazole BP

COMPOSITION

Afun[®] VT 100: Each vaginal tablet contains 100 mg Clotrimazole BP.

Afun[®] VT 200: Each vaginal tablet contains 200 mg Clotrimazole BP.

Afun[®] VT 500: Each vaginal tablet contains 500 mg Clotrimazole BP.

PHARMACOLOGY

Clotrimazole is a broad spectrum antifungal. It also exhibits activity against *Trichomonas*, *Staphylococci*, *Streptococci* and *Bacteroides*. It inhibits biosynthesis of sterols, particularly ergosterol, an essential component of the fungal cell membrane, thereby damaging and affecting the permeability of the cell membrane.

INDICATION

Vaginal itching, burning and discharge associated with recurrent vaginal yeast infections (Vaginal candidiasis) and also in vaginitis due to *Candida* species as well as Mycotic infections complicated by other microorganisms sensitive to the drug.

DOSAGE AND ADMINISTRATION

Afun[®] VT 100 & 200:

First infections: In general, a 3-day treatment is sufficient for *Candida* vaginitis. On 3 consecutive nights, 2 **Afun[®] VT 100** or 1 **Afun[®] VT 200** are inserted as deeply as possible into the vagina. This is best achieved when lying on one's back with the knees slightly bent.

Re-infections: In general, a 6-day treatment is sufficient. This applies also to mixed infections with the above-mentioned causative organisms. On six consecutive nights, 1 **Afun[®] VT 100** is inserted as deeply as possible into the vagina. If necessary, 2 **Afun[®] VT 100** (1 in the morning and 1 in the evening) or 1 **Afun[®] VT 200** daily can be used for 6-12 days. Treatment should be timed so as to avoid the menstrual period and be finished before the onset of menstruation. To avoid re-infection, the partner should undergo local treatment with **Afun[®]** cream at the same time.

Afun[®] VT 500: Single dose, complete treatment with one vaginal tablet. The tablet should be inserted as deeply as possible into the vagina. This is best achieved when lying on one's back with the knees slightly bent. It is recommended that the treatment should be timed so as to avoid the menstrual period. For prevention of re-infection the partner should be treated locally with **Afun[®]** cream at the same time.

Afun[®] vaginal tablets are colorless and do not stain the underwear.

CONTRAINDICATION AND PRECAUTION

Hypersensitivity to Clotrimazole.

SIDE EFFECT

Since there is practically no Clotrimazole absorption through the vaginal skin, no systemic effect is expected. The local tolerance of Clotrimazole vaginal tablet is generally good. Local irritation or burning sensation may occur in a very few cases but these symptoms are not considered harmful.

DRUG INTERACTION

No information is available.

USE IN PREGNANCY

It is recommended that Clotrimazole should be used in pregnancy only when considered necessary by the physician.

STORAGE

Store below 25°C. Protect from light and moisture.

HOW SUPPLIED

Afun[®] VT 100: Each box contains 6 vaginal tablets with applicator.




Afun[®] VT 200: Each box contains 3 vaginal tablets with applicator.

Afun[®] VT 500: Each box contains 1 vaginal tablet with applicator.

DIRECTION FOR USE

1. Pull out plunger 'A' until it stops. Place a vaginal tablet into the applicator 'B'
2. Insert applicator containing the tablet carefully as deeply as possible into the vagina (best lying on your back)
3. Push plunger 'A' until it stops, thereby depositing the tablet into the vagina. Remove the applicator
4. After use, remove plunger 'A' completely by pulling it out of the applicator 'B'. Then wash it in warm (not boiling) soapy water, rinse and dry carefully

Note: Pregnant women must follow physician's instructions



Manufactured by :



SQUARE
PHARMACEUTICALS LTD.
BANGLADESH

© Registered Trade Mark.

Revision No.:01