

# Asynta™

Sodium Alginate and Potassium Bicarbonate

## COMPOSITION

**Asynta™** Tablet: Each Chewable Tablet contains Sodium Alginate USP 500 mg and Potassium Bicarbonate USP 100 mg.

**Asynta™** Suspension: Each 5 ml Suspension contains Sodium Alginate USP 500 mg and Potassium Bicarbonate USP 100 mg.

## PHARMACOLOGY

**Asynta™** belongs to a group of medicines called 'Reflux Suppressants'. Reflux is the process in which the acid juices in the stomach flow back up into the food pipe causing the pain and discomfort commonly known as heartburn. **Asynta™** works by forming a protective layer that floats on the top of the stomach contents, preventing reflux and keeping the stomach contents away from the lining of the food pipe.

## INDICATION

Treatment of symptoms of gastroesophageal reflux such as acid regurgitation, heartburn and indigestion (related to reflux), for example, following meals, or during pregnancy, or in patients with symptoms related to reflux oesophagitis.

## DOSAGE AND ADMINISTRATION

**Asynta™** Tablet: Daily 4 times, after meals and at bedtime.

Adult and children over 12 years: 1-2 tablets to be chewed after meals and at bedtime.

Children 6- 12 years: 1 tablet to be chewed after meals and at bedtime.

**Asynta™** Suspension: Daily 4 times, after meals and at bedtime.

Adult and children over 12 years: 1-2 teaspoonfuls (5-10 ml) Suspension.

Children 2 - 12 years:  $\frac{1}{2}$  - 1 teaspoonfuls (2.5-5 ml) Suspension.

## CONTRAINDICATION

This medicinal product is contraindicated in patients with known or suspected hypersensitivity to the active substances or to any of the excipient.

## DRUG INTERACTION

Interaction with other medicinal products and other forms of interaction are unknown.

## PREGNANCY AND LACTATION

**Asynta™** can be given in pregnant and lactating mother.

## SIDE EFFECT

Very rarely (<1/10,000) patients may develop allergic manifestations, such as urticaria or bronchospasm, anaphylactic or anaphylactoid reactions.

## OVERDOSE

In the event of overdose symptomatic treatment should be given. The patient may notice abdominal distension.

## STORAGE

**Asynta™** Tablet: Store below 30°C, protect from light and moisture. Keep all medicines out of the reach of the children.

**Asynta™** Suspension: Store in a cool and dry place, protect from light and moisture. Keep all medicines out of the reach of the children.

## HOW SUPPLIED

**Asynta™** Tablet: Each box contains 50 tablets in blister pack.

**Asynta™** Suspension: Each PET bottle contains 200 ml of suspension.

Manufactured by



**SQUARE**  
**PHARMACEUTICALS LTD.**

Pabna, Bangladesh

TM - Trade Mark

Revision No.:00