

Cinaron®

Cinnarizine BP

Cinaron® provides fast relief of various symptoms of cerebral and peripheral circulation disorders and prevents attacks of vascular headache such as migraine. **Cinaron®** normalizes cerebral circulation time and curves and improves cerebral irrigation in patients suffering from cerebral circulation disorders. **Cinaron®** normalizes arterial pulsations in patients suffering from vasospastic disorders (Raynaud's disease). **Cinaron®** significantly increases the rate of inflow of pulsations and the blood flow in the most affected limbs as well as the walking distance of patients suffering from intermittent claudication. **Cinaron®** reduces the healing time of crural ulcers. **Cinaron®** reduces or eliminates symptoms of vestibular disturbances: vertigo, tinnitus and autonomic manifestations (nausea, vomiting, pallor and cold extremities) and reduces or prevents future episodes.

COMPOSITION

Each tablet contains Cinnarizine BP 15 mg.

INDICATIONS

1. Cerebral circulatory disorders:

- a) Prophylaxis and maintenance therapy for symptoms of cerebral vascular spasms and arteriosclerosis such as : dizziness, ear buzzing (tinnitus), vascular headache, unsociability and irritability, fatigue, sleep rhythm disorders such as premature awakening, involuntal depressions, loss of memory and lack of concentration, incontinence and other disorders due to aging.
- b) Sequelae of cerebral and cranial trauma.
- c) Postapoplectic disorders.
- d) Migraine.

2. Peripheral circulatory disorders:

Prophylaxis and maintenance therapy for symptoms of vascular spasms and arteriosclerosis (obliterating arteritis, thromboangitis obliterans, Raynaud's disease, diabetes, acrocyanosis pernio etc.) such as intermittent claudication, trophic disturbances, pregangrene, trophic and varicose ulcers, paraesthesia, nocturnal cramps, cold extremities.

3. Disorders of balance:

- a) Prophylaxis and maintenance therapy for symptoms of labyrinthine arteriosclerosis, vestibular irritability, Meniere's syndrome such as vertigo, dizziness, giddiness, syncopal attacks, tinnitus, nystagmus, nausea and vomiting.
- b) Prophylaxis of motion sickness.

DOSAGE AND ADMINISTRATION

Usual adult dose: 15 to 30 mg three times daily. *Children (5 to 12 years):* Half of the adult dose.

Motion sickness: A dose of 30 mg two hours before the start of Journey and 15 mg every 8 hours during the journey.

Peripheral arterial disease: 75 mg two or three times daily, or as prescribed by the physician.

CONTRAINDICATIONS AND SIDE EFFECTS

Contra indications are not known. Side-effects such as somnolence and gastro-intestinal disturbances are extremely rare. They are transient and may be readily prevented by achieving the optimal dosage gradually. Combination therapy with other drugs is possible whenever indicated.

USE IN PREGNANCY AND LACTATION

Pregnancy: The safety of Cinnarizine in human pregnancy has not been established.

Lactation: Cinnarizine should not be taken by women who are breast feeding.

STORAGE CONDITION

Store below 30°C, away from light & moisture. Keep Cinaron® out of the reach and sight of children.

HOW SUPPUED

Box containing 200 tablets in blister pack.

Manufactured by



SQUARE
PHARMACEUTICALS LTD.

Kaliakoir, Bangladesh

®-Registered Trade Mark