

Doxytin-Vet[®] Powder

Doxycycline Hyclate & Colistin Sulphate

Composition:

Each gram powder contains Doxycycline 100 mg and Colistin 5,00,000 IU.

Indication:

Doxytin-Vet[®] Powder is effective against various gm(+)ve & gm(-)ve bacteria such as E.Coli, Salmonella as well as Mycoplasma. Doxytin-Vet[®] Powder is used to treat & prevent –

- Salmonellosis
- Colibacillosis
- Infectious Coryza
- Chronic Respiratory Disease (CRD)
- Mycoplasmosis in poultry.

Benefits of Doxytin-Vet[®] Powder:

- Doxytin-Vet[®] Powder is mainly active against Respiratory tract & Gastrointestinal tract infections.
- Doxytin-Vet[®] Powder is poorly absorbed from the GIT which ensures longer action at GIT; it develops its main action in the intestine against E.coli, Salmonella & Enterobacter.
- Doxytin-Vet[®] Powder has good affinity towards the lungs & is therefore especially useful for the treatment of bacterial respiratory infection.
- Doxytin-Vet[®] Powder is effective at low doses.
- Doxytin-Vet[®] Powder prevents mixed infections & mortality of day old chicks.

Dosage & Administration:

Poultry:

<2 month: 0.5 gm/litre of drinking water for 3-5 days.

>2 month: 1 gm / litre of drinking water for 3-5 days.

In Feed:

For Treatment: 100-150 gm Doxytin-Vet[®] Powder mixed with 100 Kg Feed for 3-5 days.

For Prevention: 30-50 gm Doxytin-Vet[®] Powder mixed with 100 Kg Feed for 5-7 days.

Calves, Goat & sheep:

2 gm/day/15 kg body weight for 5 days.

or as prescribed by the registered veterinary doctor.

Precaution:

Don't use to animals with seriously impaired renal and/or hepatic functions. It should not be used to the animals hypersensitive to Tetracycline. Doxytin-Vet[®] Powder should not be used combine with Penicillin & Cephalosporin.

Withdrawal Period:

Poultry meat-7 days.

Pack Size:

100 gm container.