

Fona™ Plus Gel

Adapalene & Benzoyl peroxide

COMPOSITION

Each gram of **Fona™ Plus Gel** contains Adapalene BP 1 mg and Benzoyl peroxide BP 25 mg.

PHARMACOLOGY

Fona™ Plus Gel is a combination of Adapalene, a retinoid, and Benzoyl peroxide, and is indicated for the topical treatment of acne vulgaris in patients 12 years of age and older. Adapalene binds to specific retinoic acid nuclear receptors but does not bind to cytosolic receptor protein. Biochemical and pharmacological profile studies have demonstrated that Adapalene is a modulator of cellular differentiation, keratinization and inflammatory processes. Adapalene binds with specific retinoic acid nuclear receptors that normalize the differentiation of follicular epithelial cells resulting in decreased microcomedone formation. Benzoyl peroxide is an oxidizing agent with bacteriocidal and keratolytic effects.

INDICATION

Fona™ Plus Gel is indicated for the topical treatment of acne vulgaris in patients 12 years of age and older.

DOSAGE AND ADMINISTRATION

Apply a thin film of **Fona™ Plus Gel** to affected areas of the face and/or trunk once daily after washing. Use a pea-sized amount for each area of the face (e.g., forehead, chin, each cheek). Avoid the eyes, lips and mucous membranes.

Fona™ Plus Gel is not for oral, ophthalmic, or intravaginal use.

CONTRAINDICATION

Should not be administered to individuals who are hypersensitive to any of its component.

PRECAUTION

Avoid exposure to sunlight and sunlamps. Wear sunscreen when sun exposure cannot be avoided.

SIDE EFFECT

Erythema, scaling, dryness, and stinging/ burning may occur. Most commonly reported adverse events are dry skin, contact dermatitis, application site burning, application site irritation, and skin irritation.

DRUG INTERACTIONS

Concomitant topical acne therapy should be used with caution because a possible cumulative irritancy effect may occur, especially with the use of peeling, desquamating, or abrasive agents. No formal drug-drug interaction studies were conducted.

USE IN PREGNANCY AND LACTATION

There are no well-controlled trials in pregnant women treated with Adapalene and Benzoyl peroxide. Animal reproduction studies have not been conducted with the combination gel or Benzoyl peroxide. Furthermore, such studies are not always predictive of human response; therefore, this preparation should be used during pregnancy only if the potential benefit justifies the risk to the fetus. It is not known whether Adapalene or Benzoyl peroxide is excreted in human milk following use. Caution should be exercised when administered to a nursing woman.

PEDIATRIC USE

Safety and effectiveness of Adapalene and Benzoyl peroxide gel in pediatric patients under the age of 12 years have not been established.

STORAGE

Store in a cool (below 25°C) and dry place protected from light and moisture. Keep out of the reach of children. Keep the tube tightly closed after use.

HOW SUPPLIED

Fona™ Plus Gel: Each pack has a laminated tube containing 10 gm gel.

Manufactured by :



SQUARE
PHARMACEUTICALS LTD.
BANGLADESH

TM - Trade Mark

Revision No.:00