

Loratin[®]

Loratadine

Composition

Loratin[®] Tablet : Each tablet contains Loratadine USP 10 mg.

Loratin[®] Suspension : Each 5 ml suspension contains Loratadine USP 5 mg.

Pharmacology

Loratadine is a non-sedative histamine H1 receptor antagonist with antiallergic properties devoid of anticholinergic activity. It is rapidly effective and long-lasting, allowing once-a-day administration.

Indication

- * Seasonal allergic rhinitis
- * Perennial allergic rhinitis
- * Skin allergies including chronic urticaria.

Dosage and Administration

Adult and child over 6 years: One Loratin[®] 10 mg tablet or two teaspoonful (10 ml) **Loratin[®]** suspension once daily.

Children aged 2-5 years: 5 ml or one teaspoonful (5 mg) suspension once daily.

Loratadine is not recommended for children under 2 years of age. Upon several weeks administration, Loratadine does not show any decreased activity.

Contraindication

Loratadine is contraindicated in patients who have shown hypersensitivity or idiosyncrasy.

Side Effect

The incidence of adverse effects, including sedation and anticholinergic effects, observed with Loratadine was comparable to that observed with placebo. Other events like fatigue, nausea, and headache were reported rarely. Tachycardia and syncope have been reported rarely.

Use in Pregnancy & Lactation

The safe use of loratadine during pregnancy or lactation has not been established and therefore the compound should be used only if the potential benefit justifies the potential risk to fetus or infant.

Drug Interaction

When administered concurrently with alcohol, Loratadine has no potentiating effects as measured by psychomotor performance studies.

Storage Condition

Store in a cool, dry place, protected from light. Store the suspension below 30°C. Keep out of reach of children.

How Supplied

Loratin[®] Tablet : Box containing 10 x 10 tablets in blister pack.

Loratin[®] Suspension: Each PET bottle contains 60 ml suspension and a measuring cup.

Manufactured by :



© Registered Trade Mark.