



Zesup[®]

Zinc sulphate
Mineral

COMPOSITION

Zesup[®] syrup : Each 5 ml syrup contains Zinc Sulphate USP equivalent to 10 mg elemental Zinc.

INDICATION

Zesup[®] syrup is indicated for the following conditions -

- Recurrent Respiratory Tract Infections
- Diarrhoea
- Loss of appetite
- Severe growth retardation
- Deformed bone formation
- Impaired immunological response
- Acrodermatitis enteropathica
- Parakeratotic skin lesions
- Defective and delayed wound healing
- Anaemia
- Night blindness
- Mental disturbances

DOSAGE AND ADMINISTRATION

Children under 10 kg: One teaspoonful 2 times daily.

Children within 10 to 30 kg: Two teaspoonful 1-3 times daily.

Adults and children over 30 kg body weight: Four teaspoonful 1-3 times daily.

CONTRAINDICATION

It is contraindicated in patients with hypersensitivity to Zinc.

PRECAUTION AND WARNING

Concurrent administration of Zinc salt with penicillamine might diminish the effect of Penicillamine. The absorption of Zinc, although poor, may be decreased by various compounds including some foods. Chelation may occur with tetracyclines.

DRUG INTERACTION

Zinc may inhibit the absorption of concurrently administered tetracyclines, when both are being given an interval of at least 3 hours.

OVERDOSE

Zinc sulphate is corrosive in overdose. Symptoms are corrosion and inflammation of the mucous membrane of the mouth and stomach;

MINERALS

Zesup[®]

ulceration of the stomach followed by perforation may occur. Gastric lavage and emesis should be avoided. Demulcents such as milk should be given. Chelating agents such as sodium edetate may be useful.

STORAGE CONDITION

Store in a cool place. The syrup should be protected from light.

HOW SUPPLIED

Zesup[®] syrup : Bottle containing 100 ml syrup.

