

# **Esloric-Vet<sup>®</sup> Powder**

Allopurinol BP

## **Composition:**

Each gm powder contains Allopurinol BP 50 mg.

## **Pharmacology:**

Allopurinol (Esloric-Vet<sup>®</sup>) impairs the conversion of hypoxanthine to xanthine & the conversion of xanthine to uric acid leading to reduction of deposition of urate crystal.

## **Indication:**

It is indicated for the prevention and treatment of high blood uric acid level and gout. It also prevents deposition of urate crystal in kidney, serous membranes of liver, heart, air sac and joints of poultry.

## **Dosage & Administration:**

Poultry- 25 mg/kg body weight or 2.5 gm/liter water for 3-5 days, or as directed by registered veterinarian.

## **Precaution:**

Should be used in recommended dose.

## **Contraindication:**

Esloric-Vet<sup>®</sup> is not recommended if the animal shows hypersensitivity to Allopurinol.

## **Use in pregnancy and lactation:**

Not applicable

## **Withdrawal Period:**

Nil

## **Side effects:**

Skin rash, diarrhea, fever etc.

## **Drug Interaction:**

Anticoagulant: Allopurinol extends the duration of anticoagulant. Diuretic: Concomitant use of allopurinol and thiazide diuretics may increase allopurinol toxicity.

## **Storage:**

Store below 30° C temperature & protect from light & moisture. Keep out of reach of children.

## **Pack Size:**

100 gm Sachet.