

Trevox[®] D Eye Drops

Levofloxacin USP 0.5% &
Dexamethasone BP 0.1%

Sterile Solution

Trevox[®] D Eye Drops is a combination of a third-generation fluoroquinolone Levofloxacin and a potent corticosteroid Dexamethasone. Levofloxacin controls infection by directly inhibiting bacterial DNA synthesis and promotes the breakage of DNA strands by inhibiting DNA gyrase. Dexamethasone effectively controls the inflammation by inhibiting the release of inflammatory mediators.

Composition

Trevox[®] D Eye Drops: Each ml contains Levofloxacin USP 5 mg as Levofloxacin Hemihydrate USP and Dexamethasone BP 1 mg as Dexamethasone Sodium Phosphate BP.

Indications & Uses

It is indicated for prevention and treatment of inflammation, and prevention of infection associated with cataract surgery in adults.

Dosage & Administration

One drop in the affected eye 4 times a day for 7 days. if deemed necessary, a doctor can recommend another 7 days of steroid eye drops.

Contraindication

This ophthalmic solution is contraindicated in the following cases:

1. Herpes simplex keratitis, varicella and other viral disease of the cornea and conjunctiva,
2. Mycobacterial infections of the eye caused by Mycobacterium tuberculosis, Mycobacterium leprae, or Mycobacterium avium,
3. Fungal diseases of ocular structures and Untreated purulent infection of the eye.

Side Effects

The most frequently reported drug-related undesirable effects seen with Levofloxacin are discomfort, stinging or burning sensation, itching in the eye, blurred or impaired vision in the eye.

Precaution & Warning

Prolonged use of steroid may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, eye dryness and fatigue and corneal edema.

Use in Pregnancy & Lactation

This drug should not be used during pregnancy or breast-feeding.

Use in Children

Due to the lack of safety and efficacy data during this age group, **Trevox[®] D** Eye Drops is contraindicated for administration to children and adolescents younger than 18 years of age.

Storage

Store in a cool & dry place, below 25°C, protected from light. Keep the bottle tightly closed. To prevent contamination, the bottle should be discarded 28 days after first opening. Keep this medicine out of the sight and reach of children.

How Supplied

Trevox[®] D Eye Drops: Each LDPE container contains 5 ml of Levofloxacin USP 0.5% and Dexamethasone BP 0.1% of sterile solution.

Manufactured by



SQUARE
PHARMACEUTICALS PLC.

Kaliakoir, Gazipur, Bangladesh

© Registered Trade Mark