

Cinaron® Plus

Cinnarizine BP + Dimenhydrinate USP

COMPOSITION

Cinaron® Plus Tablet : Each tablet contains Cinnarizine BP 20 mg and Dimenhydrinate USP 40 mg.

PHARMACOLOGY

Cinaron® Plus consists of two compounds Cinnarizine & Dimenhydrinate. Cinnarizine provides faster relief of various symptoms of cerebral and peripheral circulation disorders and prevents attacks of vascular headache such as migraine.

Cinnarizine normalizes cerebral circulation time and curves and improves cerebral irrigation in patients suffering from cerebral circulation disorders. Cinnarizine normalizes arterial pulsations in patients suffering from vasospastic disorders (Raynaud's disease).

Cinnarizine significantly increases the rate of inflow of pulsations and the blood flow in the most affected limbs as well as the walking distance of patients suffering from intermittent claudication. Cinnarizine reduces the healing time of crural ulcers. Cinnarizine reduces or eliminates symptoms of vestibular disturbances like, vertigo, tinnitus and autonomic manifestations (nausea, vomiting, pallor and cold extremities) and reduces or prevents future episodes.

Dimenhydrinate acts as antihistamine with anti-cholinergic properties. It exhibits anti-emetic & anti-vertigo effects through influencing the chemoreceptor trigger zone.

INDICATION

1. Cerebral circulatory disorders:

- a) Prophylaxis and maintenance therapy for symptoms of cerebral vascular spasms and arteriosclerosis such as : dizziness, ear buzzing (tinnitus), vascular headache, unsociability and irritability, fatigue, sleep rhythm disorders such as premature awakening, involuntal depressions, loss of memory, lack of concentration, incontinence and other disorders due to aging.
- b) Sequelae of cerebral and cranial trauma.
- c) Postapoplectic disorders.
- d) Migraine.

2. Peripheral circulatory disorders:

Prophylaxis and maintenance therapy for symptoms of vascular spasms an arteriosclerosis (obliterating arteritis, thromboangitis obliterans, Raynaud disease, diabetes, acrocyanosis pernio. etc.) such as intermitten claudication, trophic disturbances, pregangrene, trophic and varicose ulcer paraesthesia, nocturnal cramps, cold extremities.

3. Disorders of balance:

- a) Prophylaxis and maintenance therapy for symptoms of labyrinthin arteriosclerosis; vestibular irritability; Meniere's syndrome such as vertigo dizziness, giddiness, syncopal attacks, tinnitus, nystagmus, nausea an vomiting.
- b) Prophylaxis of motion sickness.

DOSAGE AND ADMINISTRATION

Usual adult dose: 1 tablet three times daily to be taken with liquid after meal or as advised by the physician. Children (under the age of 18): No recommended.

CONTRAINDICATIONS AND SIDE EFFECT

Contraindications are not known. Side effects such as somnolence and gastro-intestinal disturbances are extremely rare. They are transient and may be readily prevented by achieving the optimal dosage gradually. Combination therapy with other drugs is possible whenever indicated.

USE IN PREGNANCY AND LACTATION

Pregnancy: The safety of the combination of Cinnarizine & Dimenhydrinate in human pregnancy has not been established.

Lactation: The combination of Cinnarizine & Dimenhydrinate is excreted in human breast milk and should not be taken by women who are breastfeeding.

STORAGE CONDITION

Store below 30°C, away from light & moisture. Keep **Cinaron®** Plus out of the reach and sight of children.

HOW SUPPLIED

Cinaron® Plus Tablet : Each box contains 100 tablets in blister pack.

Manufactured by



SQUARE
PHARMACEUTICALS LTD.
BANGLADESH