

Composition

Truxil [™] tablet: Each film-coated tablet contains Almitrine Bismesylate INN 30 mg & Raubasine INN 10 mg.

Indication

- Reduces Neurological damage & accelerates recovery after stroke
- Minor age related neurological disorders
- Visual disorders related to age
- Disorders in the inner ear (hearing loss, dizziness, buzzing etc.)

Dosage & Administration

Usually 1-2 tablets daily (with a several hours interval)

Contraindication

This medicine is contraindicated to those patients having known hypersensitivity to any component of this preparation. Without this if anybody suffers from liver disease, peripheral neuropathy or history of peripheral neuropathy, they should not take this medicine.

Drug interactions

This drug should not be used with other medicine containing Almitrine. Due to the presence of lactose, this drug should not be used in case galactosemia, glucose or galactose mal-absorption syndrome or in lactose deficiency.

Use in pregnancy & lactation

Consulting with physicians is required to take this medicine.

Side-effects

Some side effects to some people may occur. Like-

- Weight loss, numbness in leg
- Nausea, heaviness or burning in stomach
- Vomiting, constipation, diarrhea,
- Insomnia, sleepiness, restlessness, anxiety, vertigo Palpitation etc.

Storage

Store in cool & dry place protected from light & moisture.

How supplied

Each box contains 30 tablets in blister packing.

Manufactured by

