

Composition: Each gram cream contains Eflornithine Hydrochloride 139 mg as Eflornithine Hydrochloride Monohydrate INN.

Pharmacology: It is postulated that topical Eflornithine Hydrochloride irreversibly inhibits skin Ornithine Decarboxylase (ODC) activity. This enzyme is necessary in the synthesis of polyamines. Animal data indicate that inhibition of ODC inhibits cell division and synthetic functions, which affect the rate of hair growth. Eflornithine Hydrochloride 13.9% Cream has been shown to retard the rate of hair growth in non-clinical and clinical studies.

Indication: Elorim[™] Cream is indicated for the reduction of unwanted hair in any part of the body (specially, facial hair in women).

Dosage and Administration: Apply a thin layer of **Elorim™ Cream** to affected areas of the face and adjacent involved areas under the chin and rub in thoroughly. Do not wash treated area for at least 4 hours. Use twice daily at least 8 hours apart or as directed by a physician.

Contraindication: This preparation is contraindicated in patients with a history of sensitivity to any components of the preparation.

Precaution: For external use only. Transient stinging or burning may occur when applied to abraded or broken skin.

Side Effect: Acne, headache, dry skin, itching, rash, folliculitis may occasionally occur.

Drug Interaction: It is not known if **Elorim™ Cream** has any interaction with other topically applied drug products.

Pregnancy & Lactation: Pregnancy Category C. It is not known whether or not Eflornithine Hydrochloride is excreted in human milk. Caution should be exercised when this Cream is administered to a nursing woman.

Pediatric Use: The safety and effectiveness of this product have not been established in pediatric patients less than 12 years of age.

Storage: Store in a dry and cool place below 30° C, protected from light. Do not freeze. Keep out of reach of children.

How Supplied: Each pack has a laminated tube containing 30 gm Cream.

Manufactured by

