

# Eromycin®

Erythromycin USP

## DESCRIPTION

Eromycin® (Erythromycin USP) is a bacteriostatic macrolide antibiotic. But may be bactericidal in high concentrations. Although the mechanism by which Eromycin® acts in reducing the inflammatory lesions of acne vulgaris is unknown, it is presumably due to the antibiotic action of the drug.

## COMPOSITION

Eromycin® Lotion contains Erythromycin USP 3% W/V.

## INDICATION

For topical treatment of acne, pimples & bacterial skin infections susceptible to Erythromycin.

## DOSAGE AND ADMINISTRATION

Apply in morning and evening to the affected areas. Before applying thoroughly wash with warm water and soap, rinse and pat dry all areas to be treated.

Apply with fingertips or applicator. Spread the medications lightly rather than rubbing it in. Wash hands after use. Acne lesions on the face, neck, shoulders, chest and back may be treated in this manner. Additional containers may be used, if needed. Each container should be used once and discarded.

## SIDE EFFECT

Erythema, desquamation, burning sensation, eye irritation, tenderness, dryness and oily skin.

## CONTRAINDICATION

Hypersensitivity to Erythromycin or to any of the other ingredients of the lotion.

## PRECAUTION

For external use only  
Keep away from eyes, nose, mouth and other mucous membranes. Keep out of the reach of children

## USE IN PREGNANCY AND LACTATION

Safety for use during pregnancy has not been established. Use only when potential benefits outweigh the potential hazards to the fetus. Erythromycin is excreted in breast milk. Exercise caution when administering to a nursing mother.

## USE IN CHILDREN

Safety and effectiveness in children less than 12 years have not been established.

## SUPERINFECTION

Use of antibiotics (especially prolonged or repeated therapy) may result in bacterial or fungal overgrowth of non-susceptible organisms. Such overgrowth may lead to a secondary infection. Take appropriate measures if superinfections occur.

## DRUG INTERACTION

Clindamycin interacts with Erythromycin.

## STORAGE

Store at room temperature and keep away from light.

## HOW SUPPLIED

Eromycin® Lotion: Bottle containing 25 ml lotion with a reticulated cork and a protective cap.

Manufactured by



**SQUARE**  
PHARMACEUTICALS LTD.  
BANGLADESH