

# Mevin™

Mebeverine Hydrochloride BP

## Composition

**Mevin™** Tablet: Each tablet contains Mebeverine Hydrochloride BP 135 mg.

**Mevin™ 200 SR** Capsule: Each sustained release capsule contains Mebeverine Hydrochloride BP 200 mg.

## Pharmacology

Mebeverine is a musculotropic antispasmodic agent used to relieve cramps or spasms of the stomach and intestine (gut). It is particularly useful in treating Irritable Bowel Syndrome (IBS) and similar conditions. It works by relaxing the muscles and helps to restore the normal movement of the gut.

## Indication

For the symptomatic treatment of Irritable Bowel Syndrome (IBS) and other conditions include:

- Chronic irritable colon
- Spastic constipation
- Mucous colitis
- Colicky abdominal pain
- Persistent non-specific diarrhoea

## Dosage & Administration

For adults, elderly and children over 10 years:

**Mevin™** Tablet: 1 tablet 03 times daily.

**Mevin™ 200 SR** Capsule: 1 capsule 02 times daily.

**Mevin™** is most effective when taken 20 minutes before meals. After several weeks when the desired effect has been obtained, the dosage may be gradually reduced.

Missed dose: If a dose of this medicine is missed, it should be taken as soon as possible. However, if it is almost time for the next dose, then skip the missed dose and the regular dosing schedule should be maintained. Dose should not be doubled at the same time to compensate the missed dose.

## Side Effect

Generally Mebeverine is well tolerated. However, few side-effects like skin rash, urticaria and angioedema may appear.

## Precaution

Caution should be exercised in porphyria or allergic reaction to this or any other medicine of this group.

## Contraindication

Not known.

## Drug Interaction

Not known.

## Use in Pregnancy & Lactation

No teratogenicity has been shown in animal experiments. However, the usual precautions concerning the administration of any drug during pregnancy should be exercised.

Mebeverine does not excrete in the breast milk after administering the therapeutic dose.

## Use in Children

Mebeverine is not recommended in children under 10 years.

## Overdose

On theoretical grounds it may be predicted that CNS excitability will occur in case of overdose. No specific antidote is known gastric lavage and symptomatic treatment is recommended.

## Storage

Store below 30°C, protected from light. Keep out of the reach of children.

## How Supplied

**Mevin™** Tablet: Each box contains 50 tablets in blister pack.

**Mevin™ 200 SR** Capsule: Each box contains 30 capsules in blister pack.

Manufactured by



**SQUARE**  
PHARMACEUTICALS LTD.  
Bangladesh