

Hemacare™ Solution

Banana flavored hematinic with Vitamin B₁₂

Composition

Each ml solution contains Ferrous Sulphate BP 0.5 mg, Copper Sulphate BP 0.2 mg, Zinc Sulphate USP 0.2 mg, Cobalt (as Sulphate) Ph. Gr. 0.1 mg & Vitamin B₁₂ BP 0.1 mg

Description

Hemacare™ Solution is an effective & efficient combination of different ingredients of which each have their individual specific functions-

Ferrous Sulphate: Prevents iron deficiency, poor absorption and folic acid deficiency during pregnancy, Helps in synthesis of hemoglobin and thus prevents anemia

Copper Sulphate: Antifungal, Antimicrobial activity, Helps in RBC production and absorption of iron. Acts as a catalyst in the synthesis of hemoglobin.

Zinc Sulphate: Helps in production of RBC, Rehydration and in parenteral nutrition. It enhances immunity, protein digestion & an important nutrient for the skin. Zn catalyzes enzyme reactions, normal growth, bone formation, appetite control etc.

Cobalt: Helps in vitamin B₁₂ synthesis by resident bacteria in the rumen and amino acid metabolism. It also increases immunity and appetite, also maintains the nervous system.

Vitamin B₁₂: Helps in maturation of RBC, involved in metabolism, use as a cofactor in DNA synthesis, important in functioning of the nervous system via synthesis of myelin, and in the maturation of red blood cells in the bone marrow.

Indication

Poultry: Prevention and treatment of iron and folic acid deficiency, poor absorption, anemia and dermatitis. Increase metabolism, immunity, appetite, egg production and hatchability.

Livestock: Prevention and treatment of deficiency diseases of nutrients, anemia caused by worm infestations, skin & hair diseases. Increase immunity, metabolism and appetite with improved parenteral health.

Dosage & administration

For oral administration only. The bottle should be shaken well before each use.

Livestock: 1 ml/kg body weight twice daily for 5-7 days.

Poultry: 1 ml/ liter of drinking water 1-2 times daily for 3-5 days.

Or, as directed by the registered Veterinarian. Only for the use of animals.

Side effect

No side effects are observed within recommended dose.

Contraindication

Not Known

Drug interaction

Not Known

Use in pregnancy & laying

Recommended to use with caution during pregnancy & Lactation.

Withdrawal Period

Egg & Meat: 0 (zero) day.

Storage

Store at below 30⁰ C temperature and dry place, away from direct sunlight. Keep out of reach of children.

Packing

100 ml, 500 ml & 1L