

# Aftovaxpur®

Inactivated, Purified vaccine against foot & mouth disease (FMD)

## Composition

### Antigen:

Purified Inactivated Foot and Mouth Disease Virus containing three (03) serotypes appropriate to the region. Type O, Type A & Type Asia-1.

### Adjuvants:

Aluminium hydroxide and saponin

## Indications

The vaccination of cattle, buffalo, sheep and goats against **Foot and Mouth Disease**.

## Administration and dosage

Shake well before use subcutaneous route

Dose: Cattle, Buffalo: 2 ml, Subcutaneously & Sheep & Goat: 1 ml, Subcutaneously according to the following schedule:

Primary course of vaccination:

Two injections, 3 to 4 weeks apart, starting at two weeks of age for young animals from unvaccinated dams or 2.5 months of age for young animals from vaccinated dams. In case of an epidemic, the first injection should be given to all animals as early as 2 weeks of age.

Boosters: Normally, boosters should be given every 4- 6 months depending on the risk and in accordance with local legislation.

## Contradictions:

None

## Precautions

Vaccinate only healthy animals

Observe usual aseptic procedures. Once opened bottles of vaccines should be used within 24 hours provided they have been stored between +2°C and +8°C.

## Side effects

Vaccination may be followed by a small local swelling and/ or slight pyrexia both of short duration.

## Withdrawal time

Nil

## Storage

Store between +2°C and 8°C, protected from the light. Do not freeze.

## Presentation

50 ml Bottle