

Urocare™

Nitrofurantoin USP

COMPOSITION

Urocare™ 100 SR Capsule: Each sustained release capsule contains Nitrofurantoin USP 100 mg.

PHARMACOLOGY

Nitrofurantoin is an antibacterial agent specific for urinary tract infections. It is highly soluble in urine, to which it may impart a brown color. Nitrofurantoin acts usually as a bacteriostatic antimicrobial, but may be bactericidal depending on the concentration of the drug and the susceptibility of the organism. The mechanism of antimicrobial action is unique among antibacterials. Nitrofurantoin is reduced by bacterial flavoproteins to reactive intermediates, which inactivate or alter bacterial ribosomal proteins and other macromolecules. Nitrofurantoin has greater antibacterial activity in acidic environments.

INDICATION

Nitrofurantoin (**Urocare™**) is indicated only for the treatment and prevention of acute uncomplicated urinary tract infections caused by susceptible strains of *Escherichia coli* or *Staphylococcus saprophyticus*. Nitrofurantoin (**Urocare™**) is also active against several other gram negative and some gram positive organisms like *Klebsiella*, *Enterobacter*, *Enterococci*, *Staphylococcus aureus* and *Epidermidis*, *Citrobacter*, *Salmonella*, *Shigella*, and *Corynebacterium*.

DOSAGE AND ADMINISTRATION

Nitrofurantoin (**Urocare™**) should be given with food to improve drug absorption and, in some patients, tolerance.

Urocare™ 100 SR Capsule

Adults and children over 12 years: One capsule every 12 hours for seven days.

Genito-urinary surgical prophylaxis - One capsule twice daily on the day of the procedure and for next 3 days. For long-term suppressive therapy in adults, a reduction of dosage to 50-100 mg at bedtime may be adequate.

CONTRAINDICATION

Anuria, oliguria, or significant impairment of renal function (creatinine clearance under 60 ml per minute or clinically significant elevated serum creatinine) are contraindications. Nitrofurantoin is contraindicated in patients with a previous history of cholestatic jaundice/hepatic dysfunction associated with Nitrofurantoin. Nitrofurantoin is also contraindicated in those patients with known hypersensitivity to it.

PRECAUTION

Patients should be instructed to complete the full course of therapy; however, they should be advised to contact their physician if any unusual symptoms occur during

therapy. Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Patients should be advised not to use antacid preparations containing Magnesium Trisilicate while taking Nitrofurantoin.

SIDE EFFECT

The most common side effects are nausea, headache and flatulence. Other side effects are diarrhea, dyspepsia, abdominal pain, constipation, emesis, dizziness, drowsiness etc.

DRUG INTERACTION

Coadministration of Nitrofurantoin with antacids containing Magnesium Trisilicate reduce both the rate and extent of absorption. Uricosuric drugs, such as Probenecid and Sulfinpyrazone, can inhibit renal tubular secretion of Nitrofurantoin.

USE IN PREGNANCY & LACTATION

Pregnancy: Pregnancy category B. It should be used during pregnancy only if clearly needed.

Lactation: Nitrofurantoin has been detected in human breast milk in trace amounts. Because of the potential for serious adverse reactions from Nitrofurantoin in nursing infants under one month of age, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

OVERDOSE

Occasional incidents of acute overdosage of Nitrofurantoin have not resulted in any specific symptoms other than vomiting. Induction of emesis is recommended. There is no specific antidote, but a high fluid intake should be maintained to promote urinary excretion of the drug.

STORAGE

Store below 30°C in dry place. Keep away from light. Keep out of reach of children.

HOW SUPPLIED

Urocare™ 100 SR Capsule: Each box contains 20 capsules in blister pack.

Manufactured by

