

# Fusid<sup>®</sup>-Vet Bolus

Furosemide BP 2 g

## Composition

Each bolus contains Furosemide BP 2 g.

## Pharmacology

Furosemide enhances urine production by decreasing the efficiency of the sodium reabsorption process, resulting in increased water elimination from the body.

## Indications

Indicated for the treatment of ascites, udder edema, pulmonary congestion, renal failure, acute non-inflammatory tissue edema etc. It also aids in the excretion of toxins or drug overdose by promoting increased urine flow.

## Dosage & administration

Route of administration: For oral administration only.

**Cattle** : 1 bolus/ animal once daily.

**Dog** : 2-4 mg/kg body weight once to three times daily.

**Cat** : 1-2 mg/kg body weight once or twice daily.

**Poultry** : 0.5-1 mg/kg body weight.

The bolus may be administered orally directly or through mixing with required amount of feed/water by grinding the bolus.

Or as directed by the Veterinary Physicians.

## Contraindication

Contraindicated to animal hypersensitivity to furosemide.

## Side effects

Common side effects: Increased urination alkalosis, uric acid retention and may rarely produce acute gout, hyperglycemia & glycosuria.

## Withdrawal time

Meat & Milk: 2 days, Egg: Not known

## Use in pregnancy & lactation

Can be used in Pregnancy and lactating animals. However, risk & necessity should be considered during administration in pregnant & lactating animal.

## Storage

Keep away from light and wet place, store below 30<sup>0</sup> C temperature.

## Packing

Each box contains 5x4's Bolus in blister strip.

For Veterinary use only