

Coxitril-Vet[®] liquid

Toltrazuril INN 2.5%

Composition:

Each ml Coxitril-Vet[®] liquid contains Toltrazuril INN 25 mg.

Indication:

Treatment & prevention of coccidiosis in poultry due to *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunette* & *E. mitis*.

Dosage & Administration:**Treatment:**

7 mg Toltrazuril/ kg body weight per day for 2 consecutive days or 28 ml Coxitril-Vet[®] Liquid/100 kg flocks for 2 consecutive days; 24 hours continuously or 8 hours each days in drinking water. For severe infections, the treatment may be repeated 5 days later.

Prevention:

7 mg Toltrazuril /kg body weight for 2 consecutive days within 10-18 days per month up to 4 months of age as a strategic administration program.

Or, as per direction of registered veterinarian/consultant.

Side effects:

In high dose egg production drop, growth retardation & polyneuritis may occur.

Drug Interaction:

Coxitril-Vet[®] liquid is compatible with other poultry feed additives & no hazardous reactions were reported.

Contraindication:

Should not be administered in flocks with impaired hepatic and renal function. Should not be used in animals hypersensitive to active ingredient.

Storage Condition:

Store in a cool & dry place. Protect from sunlight & moisture. Keep out of reach of children.

Pack Size:

100 ml & 500 ml.

For veterinary use only