

Norvis®

Tiemonium Methylsulphate INN

COMPOSITION

Norvis® Tablet: Each film coated tablet contains Tiemonium Methylsulphate INN 50 mg. **Norvis®** Injection: Each 2 ml ampoule contains Tiemonium Methylsulphate INN 5 mg. **Norvis®** Syrup: Each 5 ml syrup contains Tiemonium Methylsulphate INN 10 mg.

PHARMACOLOGY

Tiemonium Methylsulphate is a quaternary ammonium antimuscarinic agent with peripheral effect similar to those of atropine. Tiemonium Methylsulphate strengthens calcium bonding with phospholipids and proteins thus stabilizing the cell membrane of the GI tract.

INDICATION

Tiemonium Methylsulphate is an antispasmodic drug. It reduces muscle spasms of the intestine, biliary system, uterus & urinary bladder. It is indicated for the pain in gastrointestinal, biliary, urinary and gynecological disease such as gastroenteritis, diarrhea, dysentery, biliary colic, enterocolitis, cholecystitis, colonopathies, mild cystitis, & spasmodic dysmenorrhoea.

DOSAGE AND ADMINISTRATION

Norvis® Tablet:

Recommended oral dose of Tiemonium Methylsulphate tablet is 2-6 tablets (100-300 mg) daily in divided dose.

Norvis® Injection:

Recommended injectable dose is 1 Tiemonium Methylsulphate Injection 3 times daily, through Intravenous route slowly or Intramuscular route.

Norvis® Syrup:

Children: 3 mg - 6 mg/kg body weight/day or 1.5 ml - 3 ml /kg body weight/day in 3 divided doses. Adult: 30 mg - 90 mg or 15 ml - 45 ml 3 times daily or as directed by physician.

DRUG INTERACTION

Tiemonium Methylsulphate should not be used with other drugs without prior consult of a registered physician to avoid possible drug interaction.

USE IN PREGNANCY

Tiemonium Methylsulphate may be used in pregnancy only if it is clearly needed by the assessment of risk benefit ratio.

USE IN LACTATING MOTHER

Tiemonium Methylsulphate may be used in lactating mother only if it is clearly needed by the assessment of risk benefit ratio.

SIDE EFFECT

Side effect with the use of Tiemonium Methylsulphate is very rare.

CONTRAINDICATION

Tiemonium Methylsulphate should not be used in Glucoma or where acute pain of eyeball with vision disturbance was found. This drug should also not be used in patients with the disorder of prostate or urinary bladder.

STORAGE

Norvis® Tablet : Store below 30° C temperature. Protect from light and moisture. Keep out of children's reach.

Norvis® Syrup : Store below 30° C temperature. Protect from light. Keep out of children's reach.

Norvis® Injection : Store below 30° C temperature. Protect from light. Keep out of children's reach.

HOW SUPPLIED

Norvis® Tablet: Each box contains 50 tablets in blister pack.

Norvis® Injection: Each box contains 10 ampoules in blister pack.

Norvis® Syrup (50 ml): Each box contains 50 ml syrup in PET bottle and a measuring cup

Manufactured by



SQUARE
PHARMACEUTICALS PLC.

Bangladesh