

Nasofix™

Phenylephrine Hydrochloride BP & Chlorhexidine Gluconate BP

COMPOSITION

Nasofix™ Nasal Cream: Each gram cream contains Phenylephrine Hydrochloride BP 2.5 mg & Chlorhexidine Gluconate BP 3 mg.

PHARMACOLOGY

Chlorhexidine acts as an antiseptic. It disrupts the membranes of bacteria and some viruses, potentially reducing their growth and spread within the nasal cavity. This can be beneficial if bacterial or viral infections contribute to nasal congestion. Phenylephrine hydrochloride is a decongestant. It works by stimulating alpha-adrenergic receptors located in the blood vessels of the nasal passages. This stimulation triggers the blood vessels to constrict, reducing blood flow and tissue swelling in the congested nasal lining. The combination of chlorhexidine (potentially reducing bacterial/viral load) and phenylephrine (causing vasoconstriction and reducing swelling) might offer some relief from nasal congestion associated with infections or allergies.

INDICATION

Nasofix™ Nasal Cream is indicated for the post-operative care of nasal mucosa and as an aid in the treatment of nasal vestibulitis.

DOSAGE AND ADMINISTRATION

Nasofix™ Nasal Cream is for nasal application only. Apply the cream to the nose 4 times a day or as directed by physician.

CONTRAINDICATION

Patients who have previously shown a hypersensitivity reaction to Phenylephrine or Chlorhexidine, although such reactions are extremely rare. Hypersensitivity to the active substances or to any of the excipients.

PRECAUTION & WARNING

Always read the label and follow the directions for use. Not for use by children less than 2 years of age. Not recommended for children 2 - 6 years of ages except on medical advice. It should not be used for more than 7 consecutive days. If you have heart problems, high blood pressure or are taking anti-depressant medication see your doctor before using this product.

SIDE EFFECT

Commonly causes nasal dryness and irritation in the nasal passages.

DRUG INTERACTION

It is not known that it has any interaction with other topically applied drug products.

USE IN PREGNANCY & LACTATION

Since safety of topical Phenylephrine & Chlorhexidine use in pregnant women has not been established, drugs of this class should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Since it is not known whether topical administration of Phenylephrine & Chlorhexidine can result in sufficient systemic absorption to produce detectable quantities in breast milk, a decision should be made to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

OVERDOSE

Accidental ingestion of the contents of tube is unlikely to have any adverse effects on the patient.

STORAGE

Store below 25°C. Keep all medicine out of reach of children.

HOW SUPPLIED

Nasofix™ Nasal Cream: Each pack has an aluminium tube containing 15 gm cream and an applicator.

Manufactured by-



SQUARE
PHARMACEUTICALS PLC.
Bangladesh

200003325