

Amodis-Vet[®] Bolus

Metronidazole BP 2000 mg

Composition:

Each Bolus contains Metronidazole BP 2000 mg.

Indication:

Animal: Metritis, Endometritis, Vaginitis, Trichomoniasis, Giardiasis, Gingivitis, Amoebic dysentery and calf scour. Post operative treatment after colonic surgery & inflammatory bowel disease (IBD).

Poultry: Histomoniasis, Giardiasis, Trichomoniasis etc.

Benefits:

- ✓ Amodis-Vet[®] Bolus is active against obligate anaerobic microorganisms both protozoa and bacteria.
- ✓ Has anti-inflammatory effect in the bowel.
- ✓ Amodis-Vet[®] Bolus prevents IBD by inducing non-specific immunity.
- ✓ Absorbed rapidly from the GI tract.
- ✓ Amodis-Vet[®] Bolus can be used with other antibiotics to treat mixed bacterial infections.
- ✓ Amodis-Vet[®] Bolus is compatible with other antibiotics including Penicillin, aminoglycosides and cephalosporins.

Dosage and Administration:

Large animal:

Orally : 40-50 mg / kg body weight (i.e., 1 bolus per 40-50 kg body weight) for 5-7 days.

Intra-uterine : 4 boli directly or powdered for douche per 100 kg body weight.

Poultry:

1 bolus / 3-6 litre drinking water for 4-6 days.

Or as directed by the Registered Veterinary Doctor.

Precaution:

Amodis-Vet[®] Bolus should not be used in young puppies and kittens. Amodis-Vet[®] Bolus should be avoided or used with caution, at reduced doses, in animals with kidney or liver disease.

Drug Interaction:

Amodis-Vet[®] Bolus is reported to elevate prothrombin time in animals on warfarin or other coumarin anticoagulants. Cimetidine can interfere with the liver's ability to break down Amodis-Vet[®] Bolus and the dose may have to be reduced. Phenobarbital and phenytoin may increase the metabolism of Amodis-Vet[®] Bolus.

Contraindication:

The drug should not be used in animals hypersensitive to active ingredients.

Use in Pregnancy and Lactation:

Amodis-Vet[®] Bolus should be avoided in pregnant animals, especially in the first trimester and should not be used in lactating animals.

Side Effect:

Usually salivation, vomiting, decreased appetite, gastro-intestinal upset etc may occur.

Storage:

Protect from light, store in a cool and dry place. Keep out of reach of children.

Pack Size:

5 x 4 Bolus.